arXiv:1207.3404v1 [math.CV] 14 Jul 2012

ON A SUBCLASS OF CLOSE-TO-CONVEX HARMONIC MAPPINGS

SUMIT NAGPAL AND V. RAVICHANDRAN

Abstract. Let $\mathcal{M}$ be the subclass of close-to-convex functions consisting of harmonic functions $f = h + \overline{g}$ with $g'(z) = zh'(z)$, $\operatorname{Re}(1 + zh''(z)/h'(z)) > -1/2$ for all $|z| < 1\}$. It is shown that the radius of starlikeness of the class $\mathcal{M}$ is $\sqrt{(37 - 8\sqrt{10})/27} \approx 0.658331$ and the radius of convexity of the class $\mathcal{M}$ is $2 - \sqrt{3}$. Coefficient estimates and growth result for the class $\mathcal{M}$ are also proved.

1. Introduction

Let $\mathcal{H}$ denote the class of all complex-valued harmonic functions f in the unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ normalized by $f(0) = 0 = f_z(0) - 1$. Let $\mathcal{S}_H$ be the subclass of $\mathcal{H}$ consisting of univalent and sense-preserving functions. Such functions can be written in the form $f = h + \bar{g}$, where

$$h(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad \text{and} \quad g(z) = \sum_{n=1}^{\infty} b_n z^n \tag{1.1}$$

are analytic and $|g'(z)| < |h'(z)|$ in $\mathbb{D}$. Let $\mathcal{S}_H^0 := \{f \in \mathcal{S}_H : f_{\bar{z}}(0) = 0\}$. Observe that $\mathcal{S}_H$ reduces to $\mathcal{S}$, the class of normalized univalent analytic functions, if the co-analytic part of f is zero. In 1984, Clunie and Sheil-Small (see [3]) investigated the class $\mathcal{S}_H$ as well as its geometric subclasses. Let $\mathcal{S}_H^*$, $\mathcal{K}_H$ and $\mathcal{C}_H$ be the subclasses of $\mathcal{S}_H$ mapping $\mathbb{D}$ onto starlike, convex and close-to-convex domains, respectively. Denote by $\mathcal{S}_H^{*0}$, $\mathcal{K}_H^0$ and $\mathcal{C}_H^0$, the class consisting of those functions f in $\mathcal{S}_H^*$, $\mathcal{K}_H$ and $\mathcal{C}_H$ respectively, for which $f_{\bar{z}}(0) = 0$.

In [6], Mocanu conjectured that the functions in the family

$$\mathcal{M} = \left\{ f = h + \overline{g} : g'(z) = zh'(z),\ \operatorname{Re}\left(1 + \frac{zh''(z)}{h'(z)}\right) > -\frac{1}{2} \text{ for all } z \in \mathbb{D} \right\}.$$

are univalent in $\mathbb{D}$. In [2], Bshouty and Lyzzaik proved this conjecture by showing that $\mathcal{M} \subset \mathcal{C}_H^0$. Recently, Bharanedhar and Ponnusamy [1] conjectured that $\mathcal{M} \subset \mathcal{S}_H^{*0}$. In Section 2, it is shown that this conjecture is false by giving a counter-example; in particular, the radius of starlikeness and the radius of convexity of this function are determined. By

---

2010 *Mathematics Subject Classification.* 30C45.

Key words and phrases. univalent harmonic mappings, starlike, close-to-convex.

The research work of the first author is supported by research fellowship from Council of Scientific and Industrial Research (CSIR), New Delhi.

finding the estimates for the coefficients a_n and b_n of a function in $\mathcal{M}$, the growth theorem for the class $\mathcal{M}$ is determined. The sharp radius of convexity of $\mathcal{M}$ and the bounds for the radius of starlikeness of the class $\mathcal{M}$ are also investigated.

2. A COUNTER-EXAMPLE

The technique of shear construction [4, Section 3.4] is used to produce the required harmonic mapping. Consider the conformal mapping $w = l(z) = z/(1-z)$ of the unit disk $\mathbb{D}$ onto the half-plane $\operatorname{Re} w > -1/2$. By [3, Theorem 5.3], the harmonic mapping $F = h+\bar{g}$ is univalent and convex in the direction of real axis, where $h - g = l$ and $g' = zh'$. The resulting linear system

$$h'(z) - g'(z) = l'(z) = \frac{1}{(1-z)^2}, \quad zh'(z) - g'(z) = 0$$

has the solution

$$h'(z) = \frac{1}{(1-z)^3}, \quad g'(z) = \frac{z}{(1-z)^3}.$$

Integration gives

$$h(z) = \frac{1}{2}[k(z) + l(z)], \quad g(z) = \frac{1}{2}[k(z) - l(z)],$$

where $k(z) = z/(1-z)^2$ is the Koebe function. Moreover, note that

$$\operatorname{Re}\left(1 + \frac{zh''(z)}{h'(z)}\right) = \operatorname{Re}\left(\frac{1+2z}{1-z}\right) > -\frac{1}{2}.$$

Thus, the harmonic mapping

$$F(z) = \operatorname{Re}\left(\frac{z}{(1-z)^2}\right) + i\operatorname{Im}\left(\frac{z}{1-z}\right) = z + \sum_{n=2}^{\infty}\frac{n+1}{2}z^n + \overline{\sum_{n=2}^{\infty}\frac{n-1}{2}z^n} \tag{2.1}$$

belongs to $\mathcal{M}$. This function F is used to prove the following theorem.

Theorem 2.1. $\mathcal{M} \not\subset \mathcal{S}_H^{*0}$

Proof. It will be shown that $F \notin \mathcal{S}_H^{*0}$ by proving that F maps the unit disk $\mathbb{D}$ onto the exterior of the parabola $v^2 = -(u + 1/4)$. This is shown by using an argument similar to that used by Clunie and Sheil-Small [3, Example 5.4, p. 14]. Applying the transformation $\zeta = (1+z)/(1-z) = \xi + i\eta$, $\xi > 0$, we get

$$F(z) = \frac{1}{4}\operatorname{Re}(\zeta^2 - 1) + \frac{i}{2}\operatorname{Im}\zeta = \frac{1}{4}(\xi^2 - \eta^2 - 1) + \frac{i}{2}\eta.$$

Observe first that each point z on the unit circle $(z \neq 1)$ is carried onto a point ζ on the imaginary axis so that $\xi = 0$ and

$$F(z) = -\frac{1}{4}(\eta^2 + 1) + \frac{i}{2}\eta,$$

which maps it onto the parabola $\{w = u + iv : v^2 = -(u + 1/4)\}$. Next observe that the positive real axis $\{\zeta = \xi + i\eta : \xi > 0,\ \eta = 0\}$ is mapped monotonically to the real interval $(-1/4, \infty)$. Finally, each horizontal line

$$\{\zeta = \xi + 2ic : \xi > 0,\ c \neq 0 \text{ is a real constant}\}$$

is carried univalently to the set

$$\{w = u + ic : u = \frac{1}{4}(\xi^2 - 4c^2 - 1),\ \xi > 0\}$$

which is the half-line $\{w = u + ic : u > -(c^2 + 1/4)\}$. This proves that $F(\mathbb{D}) = \{u + iv : v^2 > -(u + 1/4)\}$. The images of concentric circles inside $\mathbb{D}$ under the harmonic mapping F are shown in Figure 1. In particular, $F \notin \mathcal{S}_H^{*0}$. For instance, $z_0 = -1 - i \in F(\mathbb{D})$ but $z_0/2 \notin F(\mathbb{D})$. In fact, $z_0/2 \in \partial F(\mathbb{D})$. ∎

Theorem 2.2. *The harmonic mapping F given by* (2.1) *sends each disk $|z| < r \leq 2 - \sqrt{3}$ to a convex domain, but the image is not convex when $2 - \sqrt{3} < r < 1$.*

Proof. The radius of convexity of F is determined by employing a calculation similar to the one carried out in [4, Section 3.5]. For the purpose of computing the radius of convexity of F, it is enough to study the change of the tangent direction

$$\Psi_r(\theta) = \arg\left\{\frac{\partial}{\partial\theta} F(re^{i\theta})\right\}$$



FIGURE 1. Graph of the function $F(z) = \operatorname{Re}\left(\frac{z}{(1-z)^2}\right) + i\operatorname{Im}\left(\frac{z}{1-z}\right)$.

of the image curve as the point $z = re^{i\theta}$ moves around the circle $|z| = r$. Note that

$$\frac{\partial}{\partial\theta}F(re^{i\theta}) = A(r,\theta) + iB(r,\theta),$$

where

$$|1-z|^6 A(r,\theta) = -r[(1-6r^2+r^4)\sin\theta + r(1+r^2)\sin 2\theta]$$

and

$$|1-z|^4 B(r,\theta) = r[(1-r^2)\cos\theta - 2r].$$

The problem now reduces to finding the values of r such that the argument of the tangent vector, or equivalently

$$\tan\Psi_r(\theta) = \frac{B(r,\theta)}{A(r,\theta)} = \frac{(1-2r\cos\theta+r^2)[2r-(1-r^2)\cos\theta]}{(1-6r^2+r^4)\sin\theta + r(1+r^2)\sin 2\theta}$$

is a non-decreasing function of θ for $0 < \theta < \pi$. A lengthy calculation leads to an expression for the derivative in the form

$$[(1-6r^2+r^4) + 2r(1+r^2)u]^2(1-u^2)\frac{\partial}{\partial\theta}\tan\Psi_r(\theta) = p(r,u),$$

where $u = \cos\theta$ and

$$\begin{aligned} p(r,u) = 1 + 4r^2 - 26r^4 + 4r^6 + r^8 - 6ur(1+r^2)(1+r^4-6r^2) \\ - 12r^2u^2(1+r^2)^2 + 4ru^3(1+r^2)(1+r^4). \end{aligned}$$

Observe that the roots of $p(r,u) = 0$ in $(0,1)$ are increasing as a function of $u \in [-1,1]$. Consequently, it follows that $p(r,u) \geq 0$ for $-1 \leq u \leq 1$ if and only if

$$p(r,-1) = (1+r)^6(1-4r+r^2) \geq 0.$$

This inequality implies that $r \leq 2-\sqrt{3}$. This proves that the tangent angle $\Psi_r(\theta)$ increases monotonically with θ if $r \leq 2-\sqrt{3}$ but is not monotonic for $2-\sqrt{3} < r < 1$. ∎

Theorem 2.3. *The harmonic mapping F given by* (2.1) *sends each disk $|z| < r \leq r_0$ to a starlike domain, but the image is not starlike when $r_0 < r < 1$, where r_0 is given by*

$$r_0 = \frac{1}{3}\sqrt{\frac{1}{3}(37-8\sqrt{10})} \approx 0.658331. \tag{2.2}$$

Proof. A similar analysis carried out in [7, Example 1.1] is used to prove the assertion. To see this, one needs to study the change of the direction $\Phi_r(\theta) = \arg F(re^{i\theta})$ of the image curve as the point $z = re^{i\theta}$ moves around the circle $|z| = r$. A direct calculation gives

$$F(re^{i\theta}) = C(r,\theta) + iD(r,\theta),$$

where

$$|1-z|^4 C(r,\theta) = r[(1+r^2)\cos\theta - 2r] \quad \text{and} \quad |1-z|^2 D(r,\theta) = r\sin\theta.$$

For our assertion, it suffices to show that

$$\tan\Phi_r(\theta)=\frac{D(r,\theta)}{C(r,\theta)}=\frac{\sin\theta(1-2r\cos\theta+r^2)}{(1+r^2)\cos\theta-2r}$$

is a nondecreasing function of θ. A straightforward calculation leads to an expression for the derivative in the form

$$[(1+r^2)u-2r]^2\frac{\partial}{\partial\theta}\tan\Phi_r(\theta)=q(r,u),$$

where $u=\cos\theta$ and

$$q(r,u)=(1-r^2)^2-2ru(1+r^2)+8r^2u^2-2r(1+r^2)u^3.$$

The problem is now to find the values of the parameter r for which the polynomial $q(r,u)$ is non-negative in the whole interval $-1\leq u\leq 1$. Observe that

$$q(r,-1)=(1+r)^4>0 \quad\text{and}\quad q(r,1)=(1-r)^4>0.$$

Also, differentiation gives

$$\frac{\partial}{\partial u}q(r,u)=-2r(1+r^2)+16r^2u-6r(1+r^2)u^2,$$

showing that $q(r,u)$ has a local minimum at $u=(4r-\sqrt{-3+10r^2-3r^4})/(3(1+r^2))$ and a local maximum at $u=(4r+\sqrt{-3+10r^2-3r^4})/(3(1+r^2))$. Thus $q(r,u)\geq 0$ for $-1\leq u\leq 1$ if and only if

$$q\left(r,\frac{4r-\sqrt{-3+10r^2-3r^4}}{3(1+r^2)}\right)=\frac{1}{27(1+r^2)^2}[27-72r^2+58r^4-72r^6+27r^8 \\ +4r(3+10r^2+3r^4)\sqrt{-3+10r^2-3r^4}]\geq 0.$$

This inequality implies that $r\leq r_0$, where r_0 is given by (2.2). This proves that the angle $\Phi_r(\theta)$ increases monotonically with θ if $r\leq r_0$ and hence the harmonic mapping F sends each disk $|z|<r\leq r_0$ to a starlike domain, but the image is not starlike when $r_0<r<1$. ▮

The image of the subdisks $|z|<2-\sqrt{3}$ and $|z|<r_0$ under F is illustrated in Figure 2 where r_0 is given by (2.2).

We conclude this section by calculating the area of the image of the subdisks $\mathbb{D}_r=\{z\in\mathbb{D}:|z|<r\}$ under the harmonic function F. Suppose $0<r<1$ and let A_r be the area of the image domain $F(\mathbb{D}_r)$. Then

$$A_r=\pi\left[r^2+\sum_{n=2}^{\infty}n(|a_n|^2-|b_n|^2)r^{2n}\right]=\pi\left[r^2+\sum_{n=2}^{\infty}n^2r^{2n}\right]=\frac{\pi r^2(1+r^2)}{(1-r^2)^3},$$

where $a_n=(n+1)/2$ and $b_n=(n-1)/2$ for $n=2,3,\ldots$.

3. FURTHER RESULTS ON THE CLASS $\mathcal{M}$

Wang *et al.* obtained coefficient estimate for close-to-convex harmonic mappings. In Theorem 3.1, the coefficient estimates for function in the class $\mathcal{M}$ are obtained. Using this, the growth estimate for functions in $\mathcal{M}$ is obtained.

Theorem 3.1. *Let $f = h+\bar{g} \in \mathcal{M}$ where h and g are given by (1.1). Then $b_1 = g'(0) = 0$,*

$$|a_n| \leq \frac{n+1}{2} \quad \text{and} \quad |b_n| \leq \frac{n-1}{2}$$

for $n = 2, 3, \ldots$. Moreover, these bounds are sharp for the function F given by (2.1).

Proof. Define $p : \mathbb{D} \to \mathbb{C}$ by

$$p(z) := 1 + \frac{2}{3}\frac{zh''(z)}{h'(z)}.$$

By hypothesis, h is locally univalent in $\mathbb{D}$, so the function $p(z) = 1 + c_1 z + c_2 z^2 + \cdots$ is analytic in $\mathbb{D}$ with $\operatorname{Re} p(z) > 0$. By the well-known Caratheodory inequality, $|c_n| \leq 2$ for $n = 1, 2, \ldots$. Now write

$$3p(z)h'(z) = 3h'(z) + 2zh'(z)$$

and compare coefficients of z^n to obtain

$$2n(n+1)a_{n+1} = 3\sum_{k=1}^{n} k a_k c_{n-k+1}, \quad n = 1, 2 \ldots,$$

where $a_1 = 1$. The proof now proceeds by induction. Suppose we have proved $|a_k| \leq (k+1)/2$ for $k = 1, 2, \ldots, n$ where $n \geq 2$. Then

$$2n(n+1)|a_{n+1}| \leq 3\sum_{k=1}^{n} k|a_k||c_{n-k+1}| \leq 3\sum_{k=1}^{n} k(k+1) = n(n+1)(n+2)$$

which proves $|a_{n+1}| \leq (n+2)/2$. Regarding the bound for b_n, note that the relation $g'(z) = zh'(z)$ gives

$$(n+1)b_{n+1} = na_n, \quad n = 1, 2, \ldots$$

so that $|b_n| \leq (n-1)/2$. This completes the proof. ▌

One of the consequence of Theorem 3.1 is the sharp upper bound on the growth of a harmonic mapping in $\mathcal{M}$.

Corollary 3.2. *The sharp inequality*

$$|f(z)| \leq \frac{|z|}{(1-|z|)^2}, \quad z \in \mathbb{D},$$

holds for every function $f \in \mathcal{M}$. Equality holds for the function F given by (2.1).

Proof. Using the estimates for $|a_n|$ and $|b_n|$ from Theorem 3.1, it follows that

$$\begin{aligned}|f(z)| &\leq |z| + \sum_{n=2}^{\infty} |a_n||z|^n + \sum_{n=2}^{\infty} |b_n||z|^n \\ &\leq |z| + \frac{1}{2}\sum_{n=2}^{\infty}(n+1)|z|^n + \frac{1}{2}\sum_{n=2}^{\infty}(n-1)|z|^n \\ &= |z| + \sum_{n=2}^{\infty} n|z|^n = \frac{|z|}{(1-|z|)^2}.\end{aligned}$$ ∎

The next theorem determines the radius of convexity of the class $\mathcal{M}$.

Theorem 3.3. *The radius of convexity of the class $\mathcal{M}$ is $2-\sqrt{3}$. Moreover, the bound $2-\sqrt{3}$ is sharp.*

Proof. Let $f = h + \bar{g} \in \mathcal{M}$. Then the analytic functions $h - e^{i\varphi}g$, $0 \leq \phi < 2\pi$ are close-to-convex in $\mathbb{D}$ (see [2]). Since the radius of convexity in close-to-convex analytic mappings is $2-\sqrt{3}$, the functions $h - e^{i\varphi}g$ are convex in $|z| < 2-\sqrt{3}$. In view of [3, Theorem 5.7], it follows that f is convex in $|z| < 2-\sqrt{3}$. The function F defined by (2.1) shows that the bound $2-\sqrt{3}$ is best possible by Theorem 2.2 ∎



FIGURE 2. Images of the subdisks $|z| < 2-\sqrt{3}$ and $|z| < \frac{1}{3}\sqrt{\frac{1}{3}(37-8\sqrt{10})}$ under the mapping F.

Theorem 3.4. *If r_S is the radius of starlikeness of $\mathcal{M}$, then*

$$4\sqrt{2}-5 \leq r_S \leq \frac{1}{3}\sqrt{\frac{1}{3}(37-8\sqrt{10})}.$$

Proof. Let $f = h+\bar{g} \in \mathcal{M}$. Then the analytic functions $F_\lambda = f+\lambda g$ are close-to-convex in $\mathbb{D}$ for each $|\lambda| = 1$. Since the radius of starlikeness for close-to-convex analytic mappings is $4\sqrt{2}-5$, it follows that each F_λ is starlike in $|z| < 4\sqrt{2}-5$. By [5, Theorem 3, p. 10], f is starlike in $|z| < 4\sqrt{2}-5 \approx 0.65685$. Also, the harmonic mapping F defined by (2.2) shows that the radius of starlikeness of $\mathcal{M}$ is less than or equal to r_0, r_0 being given by (2.2). ∎

References

[1] S. V. Bharanedhar and S. Ponnusamy, Coefficient conditions for harmonic univalent mappings and hypergeometric mappings, Rocky Mountain J. Math. (2012), 18 pages, To appear.

[2] D. Bshouty and A. Lyzzaik, Close-to-convexity criteria for planar harmonic mappings, *Complex Anal. Oper. Theory* (2011) 5:767-774, DOI 10.1007/s11785-010-0056-7.

[3] J. Clunie and T. Sheil-Small, Harmonic univalent functions, *Ann. Acad. Sci. Fenn. Ser. A. I* **9** (1984), 3-25

[4] P. Duren, *Harmonic Mappings in the Plane*, Cambridge Univ. Press, 2004.

[5] R. Hernández and M. J. Martín, Stable geometric properties of analytic and harmonic functions. Preprint. Available from http://www.uam.es/mariaj.martin.

[6] P. T. Mocanu, Injectivity conditions in the complex plane, *Complex Anal. Oper. Theory* (2011) 5:759-766, DOI 10.1007/s11785-010-0052-y.

[7] Sumit Nagpal, V. Ravichandran, Fully Starlike and Convex Harmonic Mappings of order α, preprint.

[8] X.-T. Wang, X.-Q. Liang and Y.-L. Zhang, Precise coefficient estimates for close-to-convex harmonic univalent mappings, *J. Math. Anal. Appl.* **263** (2001), no. 2, 501–509.

Department of Mathematics, University of Delhi, Delhi–110 007, India

E-mail address: `sumitnagpal.dugmail.com`

Department of Mathematics, University of Delhi, Delhi–110 007, India and School of Mathematical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

E-mail address: `vravi68gmail.com`